

The critical points of a polynomial *

Zaizhao Meng

In this paper, we obtain new results on the critical points of a polynomial, these results are useful to the Sendov conjecture.

1 Introduction

Let $a \in (0, 1)$,

$$p(z) = (z - a) \prod_{k=1}^{n-1} (z - z_k), \quad |z_k| \leq 1 (k = 1, \dots, n-1)$$

with

$$p'(z) = n \prod_{j=1}^{n-1} (z - \zeta_j).$$

Let $r_k = |a - z_k|$, $\rho_j = |a - \zeta_j|$ for $k, j = 1, 2, \dots, n-1$. By relabeling we suppose that

$$\rho_1 \leq \rho_2 \leq \dots \leq \rho_{n-1}, \quad r_1 \leq r_2 \leq \dots \leq r_{n-1}. \quad (1.1)$$

We have (see [4], [5], [6])

$$2\rho_1 \sin\left(\frac{\pi}{n}\right) \leq r_k \leq 1 + a, \quad k = 1, 2, \dots, n-1. \quad (1.2)$$

Lemma A. If $0 < a < 1$ and $\rho_1 \geq 1$, then

$$|p(z)| > 1 - (1 - \lambda)^n, \quad \text{for } |z - a| = \lambda \leq \sin(\pi/n).$$

This is Lemma 2.16 of [2] (see [1]).

Theorem. If $1 - (1 - |p(0)|)^{\frac{1}{n}} \leq \lambda \leq \sin(\frac{\pi}{n})$ and $\lambda < a$, $\rho_1 \geq 1$, then there exists a critical point $\zeta_0 = a + \rho_0 e^{i\theta_0}$ such that $\Re \zeta_0 \geq \frac{1}{2}(a - \frac{\lambda(\lambda+2)}{a})$.

This theorem improves the previous known results (see [2], [3], [4]).

*2000 Mathematics Subject Classification: primary 30C15, Keywords: critical points, polynomial, derivative.

2 Proof of the Theorem

We apply Lemma A to conclude that

$$|p(z)| > 1 - (1 - \lambda)^n, \quad |z - a| = \lambda.$$

Since $p(z)$ is univalent in $|z - a| \leq \lambda$, it follows that there exists a unique point z_0 with $|z - a| \leq \lambda$ such that $p(0) = p(z_0)$. We assume that $\Im z_0 \geq 0$ (if not, consider $\overline{p(\overline{z})}$).

By a variant of the Grace-Heawood theorem, there exists a critical point in each of the half-planes bounded by the perpendicular bisector L of the segment from 0 to z_0 . Let $\zeta_0 = a + \rho_0 e^{i\theta_0}$ be the critical point in the half-plane containing z_0 .

The equation of L is $|z| = |z - z_0|$, that is

$$z\overline{z} = (z - z_0)(\overline{z} - \overline{z_0}), \quad (2.1)$$

then

$$|z_0|^2 = z\overline{z_0} + \overline{z}z_0. \quad (2.2)$$

Let $z^* = e^{i\beta_0}$ be the joint point of L and the circle $|z| = 1$, $\Im z^* \geq 0$, then

$$|z_0|^2 = e^{i\beta_0}\overline{z_0} + e^{-i\beta_0}z_0.$$

Hence

$$e^{i\beta_0} = \frac{z_0}{2} \pm \frac{z_0}{2|z_0|} \sqrt{|z_0|^2 - 4},$$

that is

$$e^{i\beta_0} = \left(\frac{1}{2} \pm \frac{i}{2|z_0|} \sqrt{4 - |z_0|^2} \right) z_0. \quad (2.3)$$

Write $z_0 = a + re^{i\alpha}$, $r \leq \lambda$, then we choose

$$\cos \beta_0 = \frac{1}{2}(a + r \cos \alpha) - \frac{1}{2|z_0|} \sqrt{4 - |z_0|^2} r \sin \alpha,$$

$$\cos \beta_0 = \frac{1}{2}(a + r \cos \alpha) - \frac{1}{2} \frac{\sqrt{4 - a^2 - 2ar \cos \alpha - r^2}}{\sqrt{a^2 + 2ar \cos \alpha + r^2}} r \sin \alpha, \quad (2.4)$$

We fix r and consider the circle $|z - a| = r$, let $\sin \alpha_1 = \frac{\sqrt{a^2 - r^2}}{a}$, $x = \cos \alpha$ and $F(x) = x - \sqrt{\frac{4 - a^2 - 2arx - r^2}{a^2 + 2arx + r^2}} \sqrt{1 - x^2}$,
then

$$\cos \beta_0 = \frac{1}{2}(a + rF(x)), \quad (2.5)$$

it is sufficient to give lower bound of $F(x)$ for $x \in [-1, -\frac{r}{a}]$.

Write $G(x) = -F(-x)$, then

$$G(x) = x + \sqrt{\frac{4 - a^2 + 2arx - r^2}{a^2 - 2arx + r^2}} \sqrt{1 - x^2}, \quad (2.6)$$

it is sufficient to give upper bound of $G(x)$ for $x \in [\frac{r}{a}, 1]$.

Write $\psi = \frac{4 - a^2 + 2arx - r^2}{a^2 - 2arx + r^2}$, $\phi = a^2 - 2arx + r^2$.

We have

$$G'(x) = 1 + \frac{1}{2}\psi^{-\frac{1}{2}}\psi'\sqrt{1 - x^2} - \psi^{\frac{1}{2}}x(1 - x^2)^{-\frac{1}{2}}, \quad (2.7)$$

and $\psi' = 8ar(a^2 - 2arx + r^2)^{-2}$,

hence

$$G'(x) = \phi^{-2}\psi^{-\frac{1}{2}}(1 - x^2)^{-\frac{1}{2}}G_1, \quad (2.8)$$

where

$$G_1 = \phi^{\frac{3}{2}}(4 - \phi)^{\frac{1}{2}}(1 - x^2)^{\frac{1}{2}} + 4ar(1 - x^2) - x\phi(4 - \phi). \quad (2.9)$$

We have $x = \frac{a^2 + r^2 - \phi}{2ar}$, then

$$G_1 = \frac{1}{2ar}G_2, \quad (2.10)$$

where

$$G_2 = \phi^{\frac{3}{2}}(4 - \phi)^{\frac{1}{2}}(4a^2r^2 - (a^2 + r^2)^2 + 2(a^2 + r^2)\phi - \phi^2)^{\frac{1}{2}} \\ + 8a^2r^2 - 2(a^2 + r^2)^2 + 4(a^2 + r^2)\phi - 2\phi^2 - (a^2 + r^2 - \phi)\phi(4 - \phi),$$

hence

$$G_2 = \phi^{\frac{3}{2}}(4 - \phi)^{\frac{1}{2}}(4a^2r^2 - (a^2 + r^2)^2 + 2(a^2 + r^2)\phi - \phi^2)^{\frac{1}{2}} \\ + 8a^2r^2 - 2(a^2 + r^2)^2 + (a^2 + r^2 + 2)\phi^2 - \phi^3,$$

and $\phi \in [(a - r)^2, a^2 - r^2]$.

The roots of G_2 satisfy

$$L = \phi^3(4 - \phi)(4a^2r^2 - (a^2 + r^2)^2 + 2(a^2 + r^2)\phi - \phi^2) = \\ (\phi^3 - (a^2 + r^2 + 2)\phi^2 + 2(a^2 + r^2)^2 - 8a^2r^2)^2 = R,$$

say.

We have

$$L = \phi^6 - 2(a^2 + r^2 + 2)\phi^5 + (8(a^2 + r^2) - 4a^2r^2 + (a^2 + r^2)^2)\phi^4 + (16a^2r^2 - 4(a^2 + r^2)^2)\phi^3, \\ R = \phi^6 - 2(a^2 + r^2 + 2)\phi^5 + (a^2 + r^2 + 2)^2\phi^4 + 4(a^2 - r^2)^2\phi^3 - 4(a^2 + r^2 + 2)(a^2 - r^2)^2\phi^2 + \\ 4(a^2 - r^2)^4.$$

Let $d = 1 - r^2$, $e = a^2 - 1$, by $L = R$, we deduce

$$ed\phi^4 - 2(e + d)^2\phi^3 + (4 + e - d)(e + d)^2\phi^2 - (e + d)^4 = 0,$$

that is

$$(e\phi^2 - 2(e + d)\phi - (e + d)^2)(d\phi^2 - 2(e + d)\phi + (e + d)^2) = 0,$$

there is only the root $\phi_0 = \frac{a^2 - r^2}{1 + r} \in [(a - r)^2, a^2 - r^2]$, and

$$x_0 = \frac{a^2 + r^2 - \phi_0}{2ar} = \frac{2r + a^2 + r^2}{2a(1 + r)} \in [\frac{r}{a}, 1].$$

We have $G'(1) < 0$, $G'(\frac{r}{a}) > 0$, $G(x_0)$ is the maxima value of $G(x)$ for $x \in [\frac{r}{a}, 1]$.

We obtain $G(x_0) = \frac{r+2}{a}$, and $G(x) \leq \frac{r+2}{a}$, $x \in [\frac{r}{a}, 1]$, hence

$$F(x) \geq -\frac{r+2}{a}, \quad x \in [-1, -\frac{r}{a}],$$

$$\cos \beta_0 \geq \frac{1}{2}(a - \frac{r(r+2)}{a}) \geq \frac{1}{2}(a - \frac{\lambda(\lambda+2)}{a}),$$

the theorem follows.

References

- [1] B.Bojanow, Q.I.Rahman, and J.Szynal, On a conjecture of Sendov about the critical points of a polynomial, Math.Z.190(1985),281-285.
- [2] I.Borcea, The sendov conjecture for polynomials with at most seven distinct zeros, Analysis 16(1996),137-159.
- [3] J.E.Brown, On the Sendov conjecture for sixth degree polynomials, Proc.Amer.Math.Soc.113(1991),939-946.
- [4] J.E.Brown and G.Xiang, Proof of the Sendov conjecture for polynomials of degree at most eight, J.Math.Anal.Appl.232(1999),272-292.
- [5] M.Marden, Conjectures on the critical points of a polynomial, Amer.Math.Monthly 90(1980),267-276.
- [6] Q.I. Rahman and G. Schmeisser, Analytic theory of polynomials, Oxford University Press, 2002 .

E-mail: mengzzh@126.com